

**Leading Causes of Death for Women  
Clark County vs. Indiana, 1995-1999**

|                 |                   | Clark County | Indiana |
|-----------------|-------------------|--------------|---------|
| All Causes      | Deaths            | 2,417        | 136,704 |
|                 | Age-Adjusted Rate | 852.23 s     | 778.40  |
| Cardiovascular  | Deaths            | 1,176        | 59,221  |
|                 | Age-Adjusted Rate | 405.09 s     | 326.70  |
| Homicide        | Deaths            | 3            | 555     |
|                 | Age-Adjusted Rate | * u          | 3.65    |
| HIV/AIDS        | Deaths            | 0            | 109     |
|                 | Age-Adjusted Rate | * u          | 0.72    |
| Lung Cancer     | Deaths            | 159          | 7,517   |
|                 | Age-Adjusted Rate | 58.62        | 45.37   |
| Breast Cancer   | Deaths            | 84           | 4,778   |
|                 | Age-Adjusted Rate | 31.16        | 29.16   |
| Ovarian Cancer  | Deaths            | 29           | 1,564   |
|                 | Age-Adjusted Rate | 10.61        | 9.48    |
| Cervical Cancer | Deaths            | 9            | 522     |
|                 | Age-Adjusted Rate | * u          | 3.31    |

s = significantly different from the state rate  
u = unstable rate (numerator less than 20)

Source: Indiana State Department of Health,  
Epidemiology Resource Center, Data Analysis Team